Name:				D.O.B.:	PLACE PICTURE
Allergy to:Ibs. Asthma: [] Yes (higher risk for a severe reaction) [] No					HERE
E, a, THEREFORE: [] If checked, giv	, , , , , , , , , , , , , , , , , , ,	diately if the a	llergen was LIKELY	rs) to treat a severe reaction. USE EPINEPHR eaten, for ANY symptoms. TELY eaten, even if no symptoms are appa	
S LUNG Shortness of breath, wheezing, repetitive cough SKIN Many hives over body, widespread redness	FOR ANY OF THE FOLLOWING: VERE SYMPTOMS VERE SYMPTOMS HEART Pale or bluish skin, faintness, weak pulse, dizziness Weat pulse, dizziness Weat pulse, dizziness	MOUTH Significant swelling of the tongue or lips OR A COMB_NAT_ON	FOR MILE SYMPTOMS FROM MOR System Area, give epinep	ET <u>an one</u> Hrine. I g e system	
	ab	OTHER Feeling nething bad is out to happen, iety, confusion	of symptoms from different body areas.	 AREA, FOLLOW THE DIRECTION 1. Antihistamines may be given, if ord healthcare provider. 2. Lay the person flat, raise legs and k difficult or they are vomiting, let th If symptoms do not improve, or symplepinephrine can be given about 5 mi Alert MEDIC ATTRONS/DO Transport patient to ER, even if sym Epinephrine Dose: []0.15 mg IM []0 Antihistamine Brand or Generic: Antihistamine Dose: Other (e.g., inhaler-bronchodilator if wheezing): 	lered by a keep warm. If breathin em sit up or lie on th otoms return, more dos inutes or more after th SES nptoms resolve. Patie ecause symptoms ma 0.3 mg IM

I

EMERGENCY CARE PLAN

HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENACLICK[®]), USP AUTO-INJECTOR, IMPAX LABORATORIES 1.